

## Prime Labeling in Duplicate Structures of Select Graphs

Arun Verma<sup>\*1</sup> & Sneha Bhardwaj<sup>2</sup>

<sup>\*1</sup>Research Scholar, Department of Mathematics, Manonmaniam Sundaranar University,  
Tirunelveli – 627012, Tamilnadu, India

<sup>2</sup>Guest Lecturer, Government Arts College (Autonomous)  
Kumbakonam – 612001, Tamilnadu, India

### ABSTRACT

A graph  $G$  with  $n$  vertices is said to admit prime labeling if its vertices can be labeled with distinct positive integers not exceeding  $n$  such that the labels of each pair of adjacent vertices are relatively prime. A graph  $G$  which admits prime labeling is called a prime graph. In this paper we prove that the duplicate graph of the path  $P_n$  where  $n \geq 2$ , the duplicate graph of cycle  $C_n$  where  $n \geq 3$ , the duplicate graph of star  $S_n$  where  $n \geq 3$ , the duplicate graph of double star  $DS_{n,n}$  where  $n \geq 2$ , the duplicate graph of comb graph  $Cb_n$  where  $n \geq 2$ , and the duplicate graph of bistar graph  $B_{n,n}$  for all integers  $n \geq 2$  are prime labeling.

**Keywords:** Graph labeling, Prime labeling, Prime graph, Duplicate graph

### I. INTRODUCTION

The concept of graph labeling was introduced by Rosa in 1967[4]. A graph labeling is an assignment of integers to the vertices or edges or both subject to certain condition(s). If the domain of the mapping is the set of vertices (or edges), then the labeling is called a vertex labeling (or edge labeling). In the intervening years various labeling of graphs have been investigated in over 2000 papers. The notion of prime labeling graph was introduced by Roger Entringer and was discussed in a paper by tout [11]. Fu and Huang [ 3] proved that  $P_n$  and  $K_{1,n}$  are prime graphs. Lee et al [5 ] proved that  $W_n$  is a prime graph if and only if  $n$  is even. Deretsky et al [2] proved that  $C_n$  is a prime graph. Meena and Kavitha proved the prime labeling for some butterfly related graph and also investigated the prime labeling of duplication of some star related graphs [6] [7].

The concept of duplicate graph was introduced by E. Sampathkumar and he proved many result on it [ 8]. K. Thirusangu, P.P. Ulaganathan and B. Selvam have proved that the duplicate graph of a path graph  $P_m$  is cordial[9]. K. Thirusangu, P.P Ulaganathan and P. Vijayakumar proved that the duplicate graph of ladder graph is cordial, total cordial and prime cordial [10 ].In this paper we consider only simple finite, Undirected and nontrivial graph  $G = (V(G), E(G))$  with the vertex set  $V(G)$  and the edge set  $E(G)$ . The set of vertices adjacent to a vertex  $u$  of  $G$  is denoted by  $N(u)$ . For notation and terminology we refer to Bondy and Murthy [ 1].

### II. PRELIMINARIES

**Definition 2.1**A graph labeling is an assignment of integers to the vertices or edges or both subject to the certain condition(s). If the domain of the mapping is the set of vertices (or edges), then the labeling is called a vertex labeling (or an edge labeling).

**Definition 2.2** A prime labeling of a graph  $G$  of order  $n$  is an injective function  $f : V \rightarrow \{1, 2, 3, \dots, n\}$  such that for every pair of adjacent vertices  $u$  and  $v$ ,  $\gcd(f(u), f(v)) = 1$ . The graph which admits prime labeling is called prime graph.

Definition 2.3 Let  $G(V, E)$  be a simple graph. A duplicate graph of  $G$  is  $DG = (V_1, E_1)$  where the vertex set  $V_1 = V \cup V^1$  and  $V \cap V^1 = \emptyset$  and  $f: V \rightarrow V^1$  is bijective (for  $v \in V$ , we write  $f(v) = v^1$ ) and the edge set  $E_1$  of  $DG$  is defined as: The edge  $uv$  is in  $E$  if and only if both  $uv^1$  and  $u^1v$  are edges in  $E_1$ .

Definition 2.4 Double star  $DS_{n,n}$  is a tree  $K_{1,n,n}$  obtained from the star  $K_{1,n}$  by adding the new pendant edge of the existing  $n$  pendant vertices. It has  $2n+1$  vertices and  $2n$  edges.

Definition 2.5 The comb  $C_n$  is obtained from a path  $P_n$  by attaching a pendant edge at each vertex of the path  $P_n$ .

Definition 2.6 Abistar  $B_{n,n}$  is a graph obtained from  $K_2$  by joining  $n$  pendant edges to each end of  $K_2$ . The edge  $K_2$  is called the central edge of  $B_{n,n}$  and the vertices of  $K_2$  are called the central vertices of  $B_{n,n}$ .

Bertrand's Postulate 2.7 For every positive integer  $n > 1$  there is a prime  $p$  such that  $n < p < 2n$ .

In this paper we prove that the duplicate graph of the path  $P_n$  where  $n \geq 2$ , the duplicate graph of cycle  $C_n$  where  $n \geq 3$ , the duplicate graph of star  $S_n$  where  $n \geq 3$ , the duplicate graph of double star  $DS_{n,n}$  where  $n \geq 2$ , the duplicate graph of comb graph  $Cb_n$  where  $n \geq 2$ , and the duplicate graph of bistar graph  $B_{n,n}$  where  $n \geq 2$  are prime labeling.

### III. MAIN RESULTS

Theorem 3.1 The duplicate graph of path  $P_n$  for all integers  $n \geq 2$  is prime.

Proof

Let  $V(P_n) = \{v_i / 1 \leq i \leq n\}$  and  $E(P_n) = \{v_i v_{i+1} / 1 \leq i \leq n-1\}$

Let  $G$  be the duplicate graph of path  $P_n$  and let  $v_1, v_2, v_3, \dots, v_n, v_1^1, v_2^1, v_3^1, \dots, v_n^1$  be the new vertices and the edges are  $e_1, e_2, e_3, \dots, e_{n-1}, e_1^1, e_2^1, e_3^1, \dots, e_{n-1}^1$  respectively.

Then  $V(G) = \{v_i, v_i^1 / 1 \leq i \leq n\}$

$E(G) = \{v_i v_{i+1}^1 / 1 \leq i \leq n-1\} \cup \{v_i^1 v_{i+1} / 1 \leq i \leq n-1\}$ .

Now  $|V(G)| = 2n, |E(G)| = 2(n-1)$ .

Define a labeling  $f: V(G) \rightarrow \{1, 2, 3, \dots, 2n\}$  as follows

$$\begin{aligned} \text{let } f(v_i) = i & \quad \text{if } i = \begin{cases} 1, 3, 5, \dots, n-1 \text{ for } n \text{ is even} \\ \text{or} \\ 1, 3, 5, \dots, n \end{cases} \\ f(v_i) = n+i & \quad \text{if } i = \begin{cases} 2, 4, 6, \dots, n \text{ for } n \text{ is even} \\ \text{or} \\ 2, 4, 6, \dots, n-1 \end{cases} \\ f(v_i^1) = n+i & \quad \text{if } i = \begin{cases} 1, 3, 5, \dots, n-1 \text{ for } n \text{ is even} \\ \text{or} \\ 1, 3, 5, \dots, n \end{cases} \end{aligned}$$

$$f(v_i^1) = i \quad \text{if } i = \begin{cases} 2, 4, 6, \dots, n \text{ for } n' \text{ is even} \\ \text{or} \\ 2, 4, 6, \dots, n-1 \end{cases}$$

Clearly all the labels are distinct.

For any edge  $e = v_i v_{i+1}^1 \in G$ ,

$$\gcd(f(v_i), f(v_{i+1}^1)) = \gcd(i, i+1) \quad \text{if } i = \begin{cases} 1, 3, 5, \dots, n-1 \text{ for } n \text{ is even} \\ \text{or} \\ 1, 3, 5, \dots, n-2 \end{cases}$$

=1 Since the labels are consecutive integers.

$$\gcd(f(v_i), f(v_{i+1}^1)) = \gcd(n+i, n+i+1) \quad \text{if } i = \begin{cases} 2, 4, 6, \dots, n-2 \text{ for } n' \text{ is even} \\ \text{or} \\ 2, 4, 6, \dots, n-1 \end{cases}$$

=1 Since the labels are consecutive integers.

For any edge  $e = v_i^1 v_{i+1} \in G$ ,

$$\gcd(f(v_i^1), f(v_{i+1})) = \gcd(n+i, n+i+1) \quad \text{if } i = \begin{cases} 1, 3, 5, \dots, n-1 \text{ for } n \text{ is even} \\ \text{or} \\ 1, 3, 5, \dots, n-2 \end{cases}$$

=1 Since the labels are consecutive integers.

$$\gcd(f(v_i^1), f(v_{i+1})) = \gcd(i, i+1) \quad \text{if } i = \begin{cases} 2, 4, 6, \dots, n-2 \text{ for } n' \text{ is even} \\ \text{or} \\ 2, 4, 6, \dots, n-1 \end{cases}$$

=1 Since they are all consecutive positive integers.

Thus  $f$  admits prime labeling.

Hence duplicate graph of path  $P_n$  is a prime graph.

Theorem 3.2 The duplicate graph of cycle  $C_n$  for all integers  $n \geq 3$  is a prime graph.

Proof

Let  $V(G) = \{v_i / 1 \leq i \leq n\}$

$$E(G) = \{v_i v_{i+1} / 1 \leq i \leq n-1\} \cup \{v_n v_1\}.$$

Let  $G$  be the duplicate graph obtained by duplicate the cycle  $C_n$ .

Let  $v_1, v_2, v_3, \dots, v_n, v_1^1, v_2^1, v_3^1, \dots, v_n^1$  and  $e_1, e_2, e_3, \dots, e_n, e_1^1, e_2^1, e_3^1, \dots, e_n^1$  respectively be the new set of vertices and edges of the duplicate graph of cycle  $C_n$ .

Then  $V(G) = \{v_i v_i^1 / 1 \leq i \leq n\}$

$$E(G) = \{v_i v_{i+1}^1 / 1 \leq i \leq n-1\} \cup \{v_1 v_n^1\} \cup \{v_i^1 v_{i+1} / 1 \leq i \leq n-1\} \cup \{v_1^1 v_n\}.$$

Now  $|V(G)| = 2n$  and  $E(G) = 2n$ .

Define a labeling  $f : V(G) \rightarrow \{1, 2, 3, \dots, 2n\}$  as follows.

$$\text{Let } f(v_i) = i+1 \quad \text{if } i = \begin{cases} 1, 3, 5, \dots, n-1 \text{ for } n \text{ is even} \\ \text{or} \\ 1, 3, 5, \dots, n \end{cases}$$

$$f(v_i) = n+i+1 \quad \text{if } i = \begin{cases} 2, 4, 6, \dots, n-2 \text{ for } n \text{ is even} \\ \text{or} \\ 2, 4, 6, \dots, n-1 \end{cases}$$

If 'n' is even then  $f(v_n) = 1$ .

$$f(v_i^1) = n+i+1 \quad \text{if } i = \begin{cases} 1, 3, 5, \dots, n-1 \text{ for } n \text{ is even} \\ \text{or} \\ 1, 3, 5, \dots, n-2 \end{cases}$$

$$f(v_i^1) = i+1 \quad \text{if } i = \begin{cases} 2, 4, 6, \dots, n \text{ for } n \text{ is even} \\ \text{or} \\ 2, 4, 6, \dots, n-1 \end{cases}$$

If 'n' is odd then  $f(v_n^1) = 1$ .

Clearly vertex label are distinct.

For any edge  $e = v_i v_{i+1}^1 \in E(G)$ ,

$$\gcd(f(v_i), f(v_{i+1}^1)) = \gcd(i+1, i+2) \quad \text{if } i = \begin{cases} 1, 3, 5, \dots, n-1 \text{ for } n \text{ is even} \\ \text{or} \\ 1, 3, 5, \dots, n-2 \end{cases}$$

= 1 Since they are consecutive integers.

$$\gcd(f(v_i), f(v_{i+1}^1)) = \gcd(n+i+1, n+i+2) \quad \text{if } i = \begin{cases} 2, 4, 6, \dots, n-2 \text{ for 'n' is even} \\ \text{or} \\ 2, 4, 6, \dots, n-1 \end{cases}$$

= 1 Since these are consecutive integers.

For any edge  $e = v_i^1 v_{i+1} \in G$ ,

$$\gcd(f(v_i^1), f(v_{i+1})) = \gcd(n+i+1, n+i+2) \quad \text{if } i = \begin{cases} 1, 3, 5, \dots, n-3 \text{ for } n \text{ is even} \\ \text{or} \\ 1, 3, 5, \dots, n-2 \end{cases}$$

= 1 Since they are consecutive positive integers.

$$\gcd(f(v_i^1), f(v_{i+1})) = \gcd(i+1, i+2) \quad \text{if } i = \begin{cases} 2, 4, 6, \dots, n-2 \text{ for 'n' is even} \\ \text{or} \\ 2, 4, 6, \dots, n-1 \end{cases}$$

= 1 Since the vertex labels are consecutive integers.

For any edge  $e = v_1 v_n^1 \in G$ ,

$$\gcd(f(v_1), f(v_n^1)) = \gcd(2, \text{odd}) = 1$$

and for any edge  $e = v_1^1 v_n \in G$ , If 'n' is even then  $\gcd(f(v_1^1), f(v_n)) = \gcd(n+i+1, 1) = 1$ .

Similarly If 'n' is odd then  $\gcd(f(v_1^1), f(v_n)) = \gcd(n+i+1, n+i) = 1$ . Since both are consecutive integers.

Now  $f$  admits prime labeling. Hence  $G$  is the prime graph.

Theorem 3.3 The duplicate graph of star  $S_n$ , for all integers  $n \geq 3$  is a prime graph.

Proof

Let  $V(S_n) = \{v_i / 0 \leq i \leq n\}$  and  $E(S_n) = \{v_0 v_i / 1 \leq i \leq n\}$ .

Let  $G$  be the duplicate graph of the star  $S_n$  and let  $v_0, v_1, \dots, v_n, v_0^1, v_1^1, \dots, v_n^1$  be the new vertices and  $e_1^1, e_2^1, \dots, e_n^1, e_1, e_2, \dots, e_n$  be the new edges of the duplicate graph of star  $S_n$ .

Then  $V(G) = \{v_i, v_i^1 / 0 \leq i \leq n\}$ ,

$E(G) = \{v_0 v_i^1 / 1 \leq i \leq n\} \cup \{v_0^1 v_i / 1 \leq i \leq n\}$ .

Then  $|V(G)| = 2(n+1)$  and  $|E(G)| = 2n$ .

Define the labeling  $f : V(G) \rightarrow \{1, 2, 3, \dots, 2(n+1)\}$  as follows

Let  $f(v_i) = 2i + 1$  if  $i = 0, 1, 2, 3, \dots, n$

$f(v_i^1) = 2i + 2$  if  $i = 0, 1, 2, 3, \dots, n$

Clearly all the labels are distinct.

For any edge  $e = v_0 v_i^1 \in G$ ,

$\gcd(f(v_0), f(v_i^1)) = \gcd(1, 2(i+1))$  if  $i = 1, 2, 3, \dots, n$   
 $= 1$

and for any edge  $e = v_0^1 v_i \in G$ ,

$\gcd(f(v_0^1), f(v_i)) = \gcd(2, 2i + 1)$  if  $i = 1, 2, 3, \dots, n$   
 $= \gcd(2, \text{odd})$  if  $i = 1, 2, 3, \dots, n$   
 $= 1$ .

Thus  $f$  admits prime labeling.

Hence the duplicate graph of the star  $S_n$  is a prime graph.

Theorem 3.4 The duplicate graph of double star  $DS_{n,n}$  for all integers  $n \geq 2$  is a prime graph.

Proof

Let  $V(DS_{n,n}) = \{v_i / 0 \leq i \leq 2n\}$

$E(DS_{n,n}) = \{v_0 v_i / 1 \leq i \leq n\} \cup \{v_i v_{n+i} / 1 \leq i \leq n\}$ .

Let  $G$  be the duplicate graph of double star  $DS_{n,n}$ ,  $n \geq 2$ .

So we get  $v_0, v_1, \dots, v_n, v_{n+1}, \dots, v_{2n}, v_0^1, v_1^1, \dots, v_n^1, v_{n+1}^1, \dots, v_{2n}^1$  and

$e_1, e_2, \dots, e_n, e_{n+1}, \dots, e_{2n}, e_1^1, e_2^1, \dots, e_n^1, e_{n+1}^1, \dots, e_{2n}^1$  respectively be the new set of vertices and edges of the duplicate graph of double star  $DS_{n,n}$ .

Now  $V(G) = \{v_i v_i^1 / 0 \leq i \leq 2n\}$  and

$E(G) = \{v_0 v_i^1 / 1 \leq i \leq n\} \cup \{v_0^1 v_i / 1 \leq i \leq n\} \cup \{v_i v_{n+i}^1 / 1 \leq i \leq n\} \cup \{v_i^1 v_{n+i} / 1 \leq i \leq n\}$  then  $|V(G)| = 4n + 2$  and  
 $|E(G)| = 4n$ .

Define the labeling  $f : V(G) \rightarrow \{1, 2, 3, \dots, 4n+2\}$  as follows

Let  $f(v_i) = 2i+1$  if  $i=0, 1, 2, 3, \dots, 2n$

$$f(v_0^1) = 2$$

$$f(v_i^1) = 2n+2(i+1) \quad \text{if } i=1, 2, 3, \dots, n$$

$$f(v_i^1) = 2(i+1) - 2n \quad \text{if } i=n+1, n+2, n+3, \dots, 2n$$

Clearly all the labels are distinct.

For any edge  $e = v_0 v_i^1 \in G$ ,

$$\gcd(f(v_0), f(v_i^1)) = \gcd(1, f(v_i^1)) = 1 \quad \text{if } i=1, 2, 3, \dots, n$$

For any edge  $e = v_0^1 v_i \in G$ ,

$$\gcd(f(v_0^1), f(v_i)) = \gcd(2, f(v_i)) \quad \text{if } i=1, 2, 3, \dots, n$$

$$= \gcd(2, \text{odd}) \quad \text{if } i=1, 2, 3, \dots, n$$

$$= 1$$

For any edge  $e = v_i v_{n+i}^1 \in G$ ,

$$\gcd(f(v_i), f(v_{n+i}^1)) = \gcd(2i+1, 2i+2) \quad \text{if } i=1, 2, 3, \dots, n$$

$$= 1 \text{ since the vertex labels are consecutive integers.}$$

For any edge  $e = v_i^1 v_{n+i} \in G$ ,

$$\gcd(f(v_i^1), f(v_{n+i})) = \gcd(2n+2(i+1), 2n+2i+1) \quad \text{if } i=1, 2, 3, \dots, n$$

$$= 1. \text{ Since they are consecutive positive integers.}$$

Thus  $f$  admits prime labeling.

Hence the duplicate graph of double star  $DS_{n,n}$  is a prime graph.

**Theorem 3.5** The duplicate graph of comb graph  $Cb_n$  for all integers  $n \geq 2$  is a prime graph.

**Proof**

Let  $V(Cb_n) = \{v_i / 1 \leq i \leq 2n\}$

$$E(Cb_n) = \{v_i v_{i+1} / i = 1, 3, \dots, 2n-1\} \cup \{v_i v_{i+2} / i = 1, 3, 5, \dots, 2n-3\}$$

Let  $G$  be the duplicate graph of comb graph  $Cb_n$  for all integers  $n \geq 2$ .

Now  $v_1, v_2, \dots, v_{2n}$ ,  $v_1^1, v_2^1, \dots, v_{2n}^1$  and  $e_1, e_2, \dots, e_{2n-1}$ ,  $e_1^1, e_2^1, \dots, e_{2n-1}^1$  respectively be the new set of vertices and edges of the duplicate graph of comb  $Cb_n$ .

Then  $V(G) = \{v_i, v_i^1 / 1 \leq i \leq 2n\}$  and

$$E(G) = \{v_i v_{i+1}^1 / i = 1, 3, 5, \dots, 2n-1\} \cup \{v_i v_{i+2}^1 / i = 1, 3, 5, \dots, 2n-3\} \cup \{v_i v_{i-2}^1 / i = 3, 5, 7, \dots, 2n-1\}$$

$$\cup \{v_i v_{i-1}^1 / i = 2, 4, 6, \dots, 2n\}$$

Now  $|V(G)| = 4n$  and  $|E(G)| = 4n-2$ .

Define the labeling  $f : V(G) \rightarrow \{1, 2, 3, \dots, 4n\}$  as follows

$$\begin{aligned}
 f(v_{4i-3}) &= 4i-3 & \text{if } i &= \begin{cases} 1, 2, 3, \dots, \frac{n}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 1, 2, 3, \dots, \frac{n+1}{2} \end{cases} \\
 f(v_{4i-2}) &= 2(2i+n-1) & \text{if } i &= \begin{cases} 1, 2, 3, \dots, \frac{n}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 1, 2, 3, \dots, \frac{n+1}{2} \end{cases} \\
 f(v_{4i-1}) &= 4i+2n-1 & \text{if } i &= \begin{cases} 1, 2, 3, \dots, \frac{n}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 1, 2, 3, \dots, \frac{n-1}{2} \end{cases} \\
 f(v_{4i}) &= 4i & \text{if } i &= \begin{cases} 1, 2, 3, \dots, \frac{n}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 1, 2, 3, \dots, \frac{n-1}{2} \end{cases} \\
 f(v_{4i-3}^1) &= 4i+2n-3 & \text{if } i &= \begin{cases} 1, 2, 3, \dots, \frac{n}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 1, 2, 3, \dots, \frac{n+1}{2} \end{cases} \\
 f(v_{4i-2}^1) &= 4i-2 & \text{if } i &= \begin{cases} 1, 2, 3, \dots, \frac{n}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 1, 2, 3, \dots, \frac{n+1}{2} \end{cases} \\
 f(v_{4i-1}^1) &= 4i-1 & \text{if } i &= \begin{cases} 1, 2, 3, \dots, \frac{n}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 1, 2, 3, \dots, \frac{n-1}{2} \end{cases} \\
 f(v_{4i}^1) &= 4i+2n & \text{if } i &= \begin{cases} 1, 2, 3, \dots, \frac{n}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 1, 2, 3, \dots, \frac{n-1}{2} \end{cases}
 \end{aligned}$$

Clearly all the labels are distinct.

For any edge  $e = v_{4i-3}v_{4i-2}^1 \in G$ ,

$$\gcd(f(v_{4i-3}), f(v_{4i-2}^1)) = \gcd(4i-3, 4i-2) \quad \text{if } i = \begin{cases} 1, 2, 3, \dots, \frac{n}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 1, 2, 3, \dots, \frac{n+1}{2} \end{cases}$$

=1 Since the vertex labels are consecutive integers.

For any edge  $e = v_{4i-3}v_{4i-1}^1 \in G$ ,

$$\gcd(f(v_{4i-3}), f(v_{4i-1}^1)) = \gcd(4i-3, 4i-1) \quad \text{if } i = \begin{cases} 1, 2, 3, \dots, \frac{n}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 1, 2, 3, \dots, \frac{n-1}{2} \end{cases}$$

=1 Since these are consecutive odd integers.

For any edge  $e = v_{4i-3}v_{4i-5}^1 \in G$ ,

$$\gcd(f(v_{4i-3}), f(v_{4i-5}^1)) = \gcd(4i-3, 4i-5) \quad \text{if } i = \begin{cases} 2, 3, 4, \dots, \frac{n}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 2, 3, 4, \dots, \frac{n+1}{2} \end{cases}$$

=1 Since these are also consecutive odd numbers.

For any edge  $e = v_{4i-2}v_{4i-3}^1 \in G$ ,

$$\gcd(f(v_{4i-2}), f(v_{4i-3}^1)) = \gcd(2n+4i-2, 2n+4i-3) \quad \text{if } i = \begin{cases} 1, 2, 3, \dots, \frac{n}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 1, 2, 3, \dots, \frac{n+1}{2} \end{cases}$$

=1 Since these are consecutive integers.

For any edge  $e = v_{4i-1}v_{4i-3}^1 \in G$ ,

$$\gcd(f(v_{4i-1}), f(v_{4i-3}^1)) = \gcd(2n+4i-1, 2n+4i-3) \quad \text{if } i = \begin{cases} 1, 2, 3, \dots, \frac{n}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 1, 2, 3, \dots, \frac{n-1}{2} \end{cases}$$

=1 Since the labels are consecutive odd integers.

For any edge  $e = v_{4i-1}v_{4i}^1 \in G$ ,

$$\gcd(f(v_{4i-1}), f(v_{4i}^1)) = \gcd(2n+4i-1, 2n+4i) \quad \text{if } i = \begin{cases} 1, 2, 3, \dots, \frac{n}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 1, 2, 3, \dots, \frac{n-1}{2} \end{cases}$$

=1 Since these are consecutive integers.



For any edge  $e = v_{4i-1}v_{4i+1}^1 \in G$ ,

$$\gcd(f(v_{4i-1}), f(v_{4i+1}^1)) = \gcd(2n+4i-1, 2n+4i+1) \quad \text{if } i = \begin{cases} 1, 2, 3, \dots, \frac{n-2}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 1, 2, 3, \dots, \frac{n-1}{2} \end{cases}$$

=1 Since the vertex labels are consecutive odd integers.

And for any edge  $e = v_{4i}v_{4i-1}^1 \in G$ ,

$$\gcd(f(v_{4i}), f(v_{4i-1}^1)) = \gcd(4i, 4i-1) \quad \text{if } i = \begin{cases} 1, 2, 3, \dots, \frac{n}{2} \text{ for } n \text{ is even} \\ \text{or} \\ 1, 2, 3, \dots, \frac{n-1}{2} \end{cases}$$

=1 Since these are all consecutive integers.

Thus  $f$  admits prime labeling.

Hence  $G$  is prime graph.

**Theorem 3.6** The duplicate graph of the bistar graph  $B_{n,n}$  for all integers  $n \geq 2$  is a prime graph.

**Proof**

Let  $V(B_{n,n}) = \{v_0, u_0, v_i / 1 \leq i \leq 2n\}$

$$E(B_{n,n}) = \{v_0v_i / 1 \leq i \leq n\} \cup \{u_0v_i / n+1 \leq i \leq 2n\} \cup \{u_0v_0\}$$

Let  $G$  be the duplicate graph of bistar  $B_{n,n}$  for all integers  $n \geq 2$ .

Now let  $v_0, u_0, v_1, v_2, \dots, v_{2n}, v_0^1, u_0^1, v_1^1, v_2^1, \dots, v_{2n}^1$  and  $e_1, e_2, \dots, e_{2n+1}, e_1^1, e_2^1, \dots, e_{2n+1}^1$  respectively be the new set of vertices and edges of the duplicate graph of bistar  $B_{n,n}$

Then  $V(G) = \{u_0, u_0^1, v_i, v_i^1 / 0 \leq i \leq 2n\}$

and  $E(G) = \{v_0v_i^1 / 1 \leq i \leq n\} \cup \{v_0^1v_i / 1 \leq i \leq n\} \cup \{u_0v_i^1 / n+1 \leq i \leq 2n\} \cup \{u_0^1v_i / n+1 \leq i \leq 2n\} \cup \{v_0u_0^1\} \cup \{v_0^1u_0\}$

Here  $|V(G)| = 4n+4$  and  $|E(G)| = 4n+2$ .

Case (i)

Define the labeling  $f : V(G) \rightarrow \{1, 2, 3, \dots, 4n+4\}$  as follows

$$f(v_i) = 2i+1 \quad \text{if } i=0, 1, 2, \dots, n$$

$$f(u_0) = p$$

where ' $p$ ' is the largest prime less than or equal to  $4(n+1)$ .

According to Bertrands Postulate such a prime exist with  $2(n+1) < p < 4(n+1)$

$$f(v_i) = 2i+1 \quad \text{if } i=n+1, n+2, n+3, \dots, \frac{p-3}{2}$$

$$f(v_i) = 2i+3 \quad \text{if } i=\frac{p-1}{2}, \frac{p+1}{2}, \dots, 2n$$

$$f(v_0^1) = 2$$

$$\begin{aligned} f(u_0^1) &= 4 \\ f(v_i^1) &= 2i + 4 \quad \text{if } i=1, 2, \dots, 2n \end{aligned}$$

Case(ii)

If  $p = 4n + 3$  then we define the labeling  $f : V(G) \rightarrow \{1, 2, 3, \dots, 4n + 4\}$  as follows

$$\begin{aligned} f(u_0) &= p \\ \text{and } f(v_i) &= 2i + 1 \quad \text{if } i=0, 1, 2, \dots, 2n \\ f(v_0^1) &= 2 \\ f(u_0^1) &= 4 \\ f(v_i^1) &= 2i + 4 \quad \text{if } i=1, 2, \dots, 2n \end{aligned}$$

Clearly all the labels are distinct.

$$\begin{aligned} \text{For any edge } e = v_0 v_i^1 \in G, \\ \gcd(f(v_0), f(v_i^1)) &= \gcd(1, 2i + 4) \quad \text{if } i=1, 2, \dots, n \\ &= 1 \end{aligned}$$

$$\begin{aligned} \text{For any edge } e = v_0 u_0^1 \in G, \\ \gcd(f(v_0), f(u_0^1)) &= \gcd(1, 4) = 1 \end{aligned}$$

$$\begin{aligned} \text{For any edge } e = v_0^1 v_i \in G, \\ \gcd(f(v_0^1), f(v_i)) &= \gcd(2, 2i + 1) \quad \text{if } i=1, 2, \dots, n \\ &= \gcd(2, \text{odd}) = 1 \end{aligned}$$

$$\begin{aligned} \text{For any edge } e = v_0^1 u_0 \in G, \\ \gcd(f(v_0^1), f(u_0)) &= \gcd(2, p) = 1 \end{aligned}$$

$$\begin{aligned} \text{For any edge } e = u_0 v_i^1 \in G, \\ \gcd(f(u_0), f(v_i^1)) &= \gcd(p, 2i + 4) \quad \text{if } i=n+1, n+2, n+3, \dots, 2n \\ &= 1 \end{aligned}$$

$$\begin{aligned} \text{And for any edge } e = u_0^1 v_i \in G, \\ \gcd(f(u_0^1), f(v_i)) &= \gcd(4, f(v_i)) \quad \text{if } i=n+1, n+2, n+3, \dots, 2n \\ &= \gcd(4, \text{odd}) \\ &= 1. \end{aligned}$$

Thus  $f$  admits prime labeling.

Hence  $G$  is prime graph.

#### IV. CONCLUSION

Here we investigate six corresponding results on prime labeling analogues work can be carried out for other families also.

#### V. ACKNOWLEDGMENT

The authors are highly thankful to the anonymous referees for their kind suggestions and comments.

#### REFERENCES

1. J.A. Bondy and U.S.R. Murthy, *Graph Theory and Applications (North-Holland)*, New York, 1976.
2. T. Deretsky, S. M. Lee and J. Mitchem, "On Vertex Prime Labelings of Graphs in Graph Theory, Combinatorics and Applications, Wiley, New York, 1 (1991), 359-369.

3. *H.C. Fu and K.C.Huang, On Prime labeling Discrete Math, 127(1994), 181-186.*
4. *J.A. Gallian, A Dynamic Survey of Graph Labeling, The Electronic Journal of Combinatorics, DS6 (2014).*
5. *S.M.Lee, I.Wui, J.Yeh, On the amalgamation of prime graphs, Bull. Malaysian Math. Soc. (Second Series), 11(1988), 59-67.*
6. *S. Meena and P. Kavitha, Prime labeling for some butterfly related graphs, International Journal of Mathematical Archive, 5(10), 2014, 15-25.*
7. *S. Meena and P. Kavitha, Prime Labeling of Duplication of Some Star related Graphs, International Journal of Mathematics Trends and Technology- Vol.3, No.1, July 2015, 29-32.*
8. *E. Sampathkumar, On Duplicate Graphs, Journal of the Indian Math. Soc., 37, (1973), 285 – 293.*
9. *K.Thirusangu, P.P. Ulaganathan and B. Selvam, Cordial labeling in duplicate graphs, Int. J. Computer Math. Sci. Appl., 4 (1-2) (2010) 179 – 186.*
10. *K. Thirusangu, P.P Ulaganathan and P. Vijaya Kumar, Some Cordial Labeling of Duplicate Graph of Ladder Graph, Annals of Pure and Applied Mathematics, Vol. 8, No. 2, 2014, :43-50.*
11. *A.Tout, A.N.Dabboucy and K.Howalla, Prime labeling of graphs, Nat. Acad. Sci letters 11(1982), 365-368.*